ASPIRIN- aspirin tablet, coated
Marc Glassman, Inc.

1018 - MAR - 2018-1206

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID [1])

---

1 nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves:
  - headache
  - muscle pain
  - toothache
  - menstrual pain
  - pain of colds
  - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose

- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, polyethylene glycol, pregelatinized starch, starch, titanium dioxide

PRINCIPAL DISPLAY PANEL

Marc’s®

NDC 68998-018-11

†Compare to the active ingredient in Bayer® Aspirin Tablets


Regular Strength

Aspirin

Pain Reliever/Fever Reducer (NSAID)

Acutal Size

100 COATED TABLETS 325 MG EACH



ASPIRIN

aspirin tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:68998-018
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients	
Ingredient Name	**Strength**
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	**Score**	no score
Shape	ROUND	**Size**	10mm
Flavor		**Imprint Code**	TCL;011
Contains			

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:68998-018-11	1 in 1 CARTON	11/01/2009	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:68998-018-14	1 in 1 CARTON	11/01/2009	10/30/2027
2		300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:68998-018-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/10/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/01/2009	

Labeler - Marc Glassman, Inc. (094487477)

Revised: 10/2026 Marc Glassman, Inc.